




# Abnormal liver function tests

Liver function tests (LFTs) are the third most common pathology test ordered by Australian GPs. Abnormal LFTs are a common finding in asymptomatic patients and can set off a cascade of further investigations. GP registrars need to have a systematic approach to interpretation of LFTs, and develop a rational approach to subsequent investigation.

<b>TEACHING AND LEARNING AREAS</b> 	<ul style="list-style-type: none"> <li>• <a href="#">Anatomy and function of the liver</a> (30 min video tutorial)</li> <li>• Risk factors for liver disease and indications for testing</li> <li>• Epidemiology and natural history of common liver diseases in general practice</li> <li>• Interpretation of abnormal LFTs (hepatocellular versus cholestatic, magnitude, rate of change)</li> <li>• Non-hepatic causes of abnormal LFTs</li> <li>• <a href="#">Investigation of abnormal LFTs</a>, including <a href="#">first and second tier testing and follow up times</a></li> <li>• Indications for referral and local pathways</li> </ul>						
<b>PRE-SESSION ACTIVITIES</b>	<ul style="list-style-type: none"> <li>• <a href="#">Read the 2011 AFP article Liver Function Tests</a></li> </ul>						
<b>TEACHING TIPS AND TRAPS</b> 	<ul style="list-style-type: none"> <li>• LFTs are a misnomer – they measure hepatocyte integrity and/or cholestasis rather than true hepatic function (except for bilirubin and albumin)</li> <li>• LFTs are not indicated for asymptomatic patients without risk factors – there is no evidence to support opportunistic testing</li> <li>• As with other tests, the ‘normal’ range of +/- 2SDs means that up to 2.5% of normal patients will have “abnormal” LFTs</li> <li>• Levels of abnormal LFTs are not a good indicator of disease severity – in fact, normal LFTs do not necessarily mean absence of liver disease</li> <li>• AST is less specific than ALT for liver disease as elevation is also caused due to damage to skeletal muscle, kidney, brain and RBCs – if only AST is elevated, consider non-liver causes</li> <li>• Consider non-liver causes if only SAP is elevated e.g. bone, placenta</li> <li>• It is vital to take a careful medication history, including OTC and herbal medications</li> <li>• An increase in the AST/ALT ratio suggests alcoholic liver disease or cirrhosis</li> <li>• Very high levels of transaminases suggest ischaemic or toxic liver injury, or viral hepatitis</li> <li>• Family history may identify risk of genetic causes of liver disease</li> <li>• Routine monitoring of LFTs is not required for patients on statins</li> </ul>						
<b>RESOURCES</b> 	<table border="1"> <tbody> <tr> <td data-bbox="336 1693 432 1794"><b>Read</b></td> <td data-bbox="432 1693 1490 1794"> <ul style="list-style-type: none"> <li>• 2005 CMAJ article <a href="#">Liver enzyme alteration: a guide for clinicians</a> – very comprehensive</li> <li>• BPAC NZ <a href="#">Liver Function Testing in Primary Care</a> – excellent summary</li> </ul> </td> </tr> <tr> <td data-bbox="336 1794 432 1850"><b>Watch</b></td> <td data-bbox="432 1794 1490 1850"> <ul style="list-style-type: none"> <li>• <a href="#">Interpreting LFTs - YouTube</a></li> </ul> </td> </tr> <tr> <td data-bbox="336 1850 432 1984"><b>Use</b></td> <td data-bbox="432 1850 1490 1984"> <ul style="list-style-type: none"> <li>• <a href="#">Device-specific checklists to assess patient’s inhaler technique</a></li> <li>• <a href="#">Lung Foundation COPD Action Plan</a></li> <li>• <a href="#">Primary Care Respiratory Toolkit – online decision support</a></li> </ul> </td> </tr> </tbody> </table>	<b>Read</b>	<ul style="list-style-type: none"> <li>• 2005 CMAJ article <a href="#">Liver enzyme alteration: a guide for clinicians</a> – very comprehensive</li> <li>• BPAC NZ <a href="#">Liver Function Testing in Primary Care</a> – excellent summary</li> </ul>	<b>Watch</b>	<ul style="list-style-type: none"> <li>• <a href="#">Interpreting LFTs - YouTube</a></li> </ul>	<b>Use</b>	<ul style="list-style-type: none"> <li>• <a href="#">Device-specific checklists to assess patient’s inhaler technique</a></li> <li>• <a href="#">Lung Foundation COPD Action Plan</a></li> <li>• <a href="#">Primary Care Respiratory Toolkit – online decision support</a></li> </ul>
<b>Read</b>	<ul style="list-style-type: none"> <li>• 2005 CMAJ article <a href="#">Liver enzyme alteration: a guide for clinicians</a> – very comprehensive</li> <li>• BPAC NZ <a href="#">Liver Function Testing in Primary Care</a> – excellent summary</li> </ul>						
<b>Watch</b>	<ul style="list-style-type: none"> <li>• <a href="#">Interpreting LFTs - YouTube</a></li> </ul>						
<b>Use</b>	<ul style="list-style-type: none"> <li>• <a href="#">Device-specific checklists to assess patient’s inhaler technique</a></li> <li>• <a href="#">Lung Foundation COPD Action Plan</a></li> <li>• <a href="#">Primary Care Respiratory Toolkit – online decision support</a></li> </ul>						
<b>FOLLOW UP/ EXTENSION ACTIVITIES</b>	<ul style="list-style-type: none"> <li>• Registrar to undertake clinical reasoning challenge and discuss with supervisor</li> </ul>						

# Abnormal liver function tests

## Clinical Reasoning Challenge

Sharon, a 51-year-old mother of four, is recalled to the surgery with abnormal LFTs (see below). She presented a week previously with tiredness and occasional itch. Sharon is otherwise totally asymptomatic. She has coeliac disease but no other significant PMHx. Other blood tests (FBC, EUC, BSL, TSH, ESR, iron studies and lipids) were normal, and LFT from 18 months ago was also normal.

### Biochemistry

Total Bilirubin	H 33	umol/L	( 4 - 20 )
Alk Phos	H 188	U/L	( 35 - 110 )
Gamma GT	H 151	U/L	( 5 - 50 )
LDH	209	U/L	( 120 - 250 )
AST	39	U/L	( 10 - 40 )
ALT	37	U/L	( 5 - 40 )
Total Protein	70	g/L	( 64 - 83 )
Albumin	47	g/L	( 36 - 47 )
Globulin	23	g/L	( 23 - 39 )

QUESTION 1. What additional key features of history are most useful in determining the cause of her abnormal LFTs?

List up to THREE.

- 1 \_\_\_\_\_
- 2 \_\_\_\_\_
- 3 \_\_\_\_\_

QUESTION 2. The remainder of her history is unremarkable. What are the MOST LIKELY diagnoses? List as many as appropriate.

---



---



---



---

QUESTION 3. What investigations would you order at this point? List as many as appropriate

---



---



---



---

# Abnormal liver function tests

## ANSWERS

### QUESTION 1

What additional key features of history are most useful in determining the cause of her abnormal LFTs? List up to THREE.

- Alcohol
- Medications (including OTC and herbal)
- Family history of liver disease

### QUESTION 2

The remainder of her history is unremarkable. What are the MOST LIKELY diagnoses? List as many as appropriate.

- Gall stones
- Fatty liver
- Malignancy e.g. pancreatic cancer, cholangiocarcinoma
- Primary biliary cirrhosis
- Primary sclerosing cholangitis

### QUESTION 3

What investigations would you order at this point? List as many as appropriate.

- Upper abdominal ultrasound
- Hepatitis B and C serology
- Autoantibodies (ANA, anti-mitochondrial antibody, anti-smooth muscle antibody)