

# Transient Ischaemic Attack (TIA) and Stroke

Transient ischaemic attacks (TIAs) are uncommon presentations in general practice and can be difficult to diagnose. However, it is essential that registrars are competent in the assessment of symptoms suggestive of a TIA, as early intervention is critical to reduce the risk of stroke. Stroke affects 50,000 Australians each year, 30 per cent of whom are under 60 years of age. Evidence suggests adherence to guidelines for management of TIA is poor. As a high-risk presentation, supervisors need to ensure that registrars are competent in assessment of patients presenting with a possible TIA or stroke.

<p><b>TEACHING AND LEARNING AREAS</b></p> 	<ul style="list-style-type: none"> <li>• Clinical features of TIAs, including typical and atypical presentations (posterior circulation)</li> <li>• Differential diagnoses and TIA mimics</li> <li>• How to undertake a neurological assessment of the patient with possible TIA</li> <li>• Immediate investigations for TIA (blood tests, brain and carotid imaging, ECG)</li> <li>• Immediate management of TIA</li> <li>• Mechanism for referral e.g. rapid access TIA clinic, ED etc</li> <li>• Screening for stroke risk</li> </ul>
<p><b>PRE-SESSION ACTIVITIES</b></p> 	<ul style="list-style-type: none"> <li>• Read the 2012 AAFP article <a href="#">TIA - diagnosis and evaluation</a></li> </ul>
<p><b>TEACHING TIPS AND TRAPS</b></p> 	<ul style="list-style-type: none"> <li>• Ensure practice staff use the <a href="#">FAST</a> mnemonic to triage patients with possible TIA/CVA</li> <li>• ~40% of patients with a diagnosis of TIA have evidence of infarction on diffusion weighted imaging</li> <li>• The average duration of TIA is approximately 10 minutes and most resolve within one hour</li> <li>• TIA is a medical emergency and precedes stroke in about 20% cases</li> <li>• Taking a very careful history is critical in the assessment of possible TIA</li> <li>• Regard all possible TIAs as high risk – use of the ABCD2 score in isolation to determine the urgency of investigation may delay recognition of AF and carotid stenosis and should be avoided</li> <li>• Urgent brain imaging is critical in order to institute antiplatelet therapy</li> <li>• Dual antiplatelet therapy should be given to patients with TIA or ischaemic stroke for 3 weeks minimum</li> <li>• Early management strategies for TIA can reduce the relative risk of stroke by 80 per cent</li> </ul>
<p><b>RESOURCES</b></p> 	<p><b>Read</b></p> <ul style="list-style-type: none"> <li>• <a href="#">Stroke Foundation Clinical Guidelines</a></li> </ul>
<p><b>FOLLOW UP/EXTENSION ACTIVITIES</b></p>	<ul style="list-style-type: none"> <li>• Ask the registrar to complete the role play under exam conditions</li> </ul>

# Transient Ischaemic Attack (TIA) and Stroke

## Role Play

### INSTRUCTIONS FOR SUPERVISOR

You are Fred Rathmines, a 69-year-old retired university lecturer, who has come to the doctor because last night you noticed that your speech was slurred.

You have been attending the practice for about seven years since moving to the area but have not met the registrar before.

*"I had some slurred speech last night and thought I should get my blood pressure checked"*

#### Story

- The episode of slurred speech came on suddenly while having dinner. You felt like you could not speak properly but had no problems thinking of what words to use. Your wife wanted to call the ambulance but you decided to wait a while and the slurred speech settled completely after about 25 minutes
- There were no other features at all i.e. weakness, visual disturbance, CVS symptoms etc.
- Nothing like this has ever happened before
- You started tablets for 'mild blood pressure' about two weeks before and despite not having any side effects so far, you wonder whether it was related to these
- You currently feel completely well, but thought that you should have a check of your BP
- You are not overly concerned about the event
- You have no significant medical problems apart from blood pressure and gout
- You had back surgery 15 years previously
- You only take the new BP medicine (ramipril) and Panadol Osteo occasionally for your back
- You don't smoke or drink
- You are married with grown up children
- You have had a TIA.

- The registrar is expected to elicit the important aspects of history, and institute an immediate management plan. When asked about examination, you can state that BP is 138/82, HR 72, BMI 28, BSL 4.1, ECG normal SR, and the remainder of the examination is completely normal.

#### Assess

- Communication skills  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
- History taking – comprehensive  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
- Problem definition – TIA likely, explanation of problem  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
- Management – urgent brain imaging, carotid dopplers, medication, urgent referral  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_
- Follow-up and safety netting  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

